

ISSN 1061-8503


ARGGIA

THE NEWS JOURNAL OF THE DRAGONFLY SOCIETY OF THE AMERICAS

VOLUME 11

31 DECEMBER 1999

NUMBER 4



Libellula quadrimaculata: the Millenium Bug

PUBLISHED BY THE DRAGONFLY SOCIETY OF THE AMERICAS

THE DRAGONFLY SOCIETY OF THE AMERICAS

Business address: c/o T. Donnelly, 2091 Partridge Lane, Binghamton NY 13903

EXECUTIVE COUNCIL 1999-2001

President	M. May	Cranbury NJ
President Elect	D. Paulson	Seattle WA
Past President	K. Tennessen	Florence AL
Past President	G.L. Harp	Jonesboro AR
Past President	T.W. Donnelly	Binghamton NY
Past President	C. Cook	Center KY
Vice President, SIO Affairs	M. Westfall, Jr.	Gainesville FL
Vice President, Canada	R. Cannings	Victoria, British Columbia
Vice President, Latin America	R. Novelo G.	Jalapa, Veracruz
Secretary	S. Dunkle	Plano TX
Treasurer	J. Daigle	Tallahassee FL
Regular member	R. Beckemeyer	Wichita KA
Regular member	J. Abbott	Denton TX
Regular member	S. Valley	Albany OR

JOURNALS PUBLISHED BY THE SOCIETY

ARGIA, the quarterly news journal of the **DSA**, is devoted to non-technical papers and news items relating to nearly every aspect of the study of Odonata and the people who are interested in them. The editor especially welcomes reports of studies in progress, news of forthcoming meetings, commentaries on species, habitat conservation, noteworthy occurrences, personal news items, accounts of meetings and collecting trips, and reviews of technical and non-technical publications. Articles for publication in **ARGIA** should preferably be submitted as hard copy and (if over 500 words) also on floppy disk (3.5" or 5.25"). The editor prefers MS DOS based files, preferably written in WORD, WORD for WINDOWS, WordPerfect, or WordStar. Macintosh WORD disks can be handled. All files should be submitted **unformatted and without paragraph indents**. Each submission should be accompanied by a text (=ASCII) file. Other languages should be submitted only as text (=ASCII) files. Line drawings are acceptable as illustrations.

T. Donnelly (address below) is the interim editor of **ARGIA**.

BULLETIN OF AMERICAN ODONATOLOGY is devoted to studies of Odonata of the New World. This journal considers a wide range of topics for publication, including faunal synopses, behavioral studies, ecological studies, etc. The **BAO** publishes taxonomic studies but will not consider the publication of new names at any taxonomic level. Enquiries and submission of manuscripts should be made to **BAO** editor T. Donnelly, 2091 Partridge Lane, Binghamton NY 13903. Final submissions (after review) should be made on floppy disk, as above, with illustrations in final form and preferably adjusted to final size.

MEMBERSHIP IN THE DRAGONFLY SOCIETY OF THE AMERICAS

Membership in the **DSA** is open to any person in any country. Dues for individuals in the US, Canada, or Latin America are \$15 for regular membership and \$20 for contributing membership, payable annually on or before 1 March of membership year. Dues for members in the Old World are \$20. **ARGIA** is mailed Air Mail outside of the US and Mexico, and First Class in those countries.

The **BULLETIN OF AMERICAN ODONATOLOGY** is available by a separate subscription at \$15 for members and \$18.75 for non-members and institutions.

Cover: *Libellula quadrimaculata*, selected as the **Millenium Bug**. It was the first species of odonate described by Linnaeus (1758) and is the subject of an article on DNA phylogeny in this issue. Drawing by Paul-Michael Brunelle

ARGIA - The News Journal of the D.S.A.

IN THIS ISSUE

Writing the date on the cover of this issue conjures up a wide range of feelings. I wonder how long it will be before I can write "2000" on checks and letters? My method of assigning locality numbers for my notebook has been to start with the two final digits of the year, thus: "99x043". "00x001" will somehow seem faintly silly.

Right now "00x001" seems far in the distance. It is very cold now, after a mild Fall (with numerous reports of late-occurring *Sympetrum vicinum*, etc.). Spring will come, and with it our next season. We should all start planning to attend the Vancouver Island DSA meeting in July. This trip will involve a lot of travel for most of our members, but they will be rewarded with a gorgeous venue with interesting Northwestern species, and a post-meeting field trip to southeastern British Columbia.

The recent passing of Juanda Bick brings a remarkable era to a close. Each of us will dwell on our personal memories of this remarkable woman.

Jerrell Daigle and his friends describe a most successful dragonfly trip to Arizona. Those who have not visited this beautiful and remarkable state should plan to do so - probably in the monsoon season in Arizona. Most of our recent new US records for odonate species have come from this state.

José Ramos fills us in with several more provincial records for odonates in the Dominican Republic. Steve Krotzer and his associates describe the recent record for *Erythemis vesiculosa* in Alabama. So-called "tropical" odonates are being captured more and more frequently in the United States.

Our main feature in this issue is Thomas Artiss' account of his DNA studies of the *Libellula* complex. Many of us have heard of Kambhampati and Charlton's results (1998) in this complex; Artiss provides a clear picture of the present state of our knowledge in this group of species. This article provides a perfect segue to my next account of the history of Odonata study in North America. Clarence Kennedy is famous for many things, but among these were his many attempts to unravel the phylogeny of various Odonata groups. I could not resist reproducing his diagram of proposed

relationships among *Libellula* species, so that readers can compare for themselves the modern DNA results and his old-fashioned techniques.

Our most important short news item is the award to Carl Cook from the StaTe of Kentucky, recognizing his many years of vigorous efforts to protect wetland habitats in that state. Well deserved, Carl!


=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=

VANCOUVER ISLAND DSA ANNUAL MEETING (WITH POSSIBLE OKANAGAN SIDE TRIP)

Gordon Hutchings , sea-trek@islandnet.com
250-995-1413

The first Annual DSA meeting of the Y2K will be held in British Columbia, the furthest western province of Canada. We will meet in Nanaimo which is a strategically located large town on Vancouver Island with adequate access via ferries and flights from the mainland, as well as an easy 1 1/2 hour drive north from Victoria, the provincial capital of tourist fame. International flights can arrive into Seattle, Vancouver as well as Victoria. Rental cars are available in these cities as well as in Nanaimo itself. We will check in Thursday evening, July 27 at 6:00 pm at the Malaspina College housing facilities and make sure most people have arrived and distribute maps. At that time, we will arrange to meet the following morning at 10:00 in a central parking lot on campus, to get out and collect. Some of the species to be found locally for the next 3 days will be *Somatochlora semicircularis*, *S. walshi*, *S. albicincta*, *Ophiogomphus occidentis*, *Cordulegaster dorsalis*, *Aeshna canadensis*, *A. multicolor*, *A. palmata*, *A. sitchensis*, *A. tuberculifera*, *Leucorrhinia glacialis*, *L. hudsonica*, *L. intacta*, *L. proxima*, *Libellula forensis*, *L. julia*, *L. lydia*, *Sympetrum illotum*, *S. madidum*, *S. obtrusum*, and *Ischnura erratica*. Chris Carson, Syd Cannings, Rob Cannings and myself will be available to lead one or more of the groups. Please bring appropriate clothing, liquids and packed lunch for all trips.

Friday, July 28: We will visit the Nanaimo Lakes district which is rich in wetland diversity ranging



George and Juanda Bick and their daughters Susan and Patty, Put-in-Bay, Ohio, 1952



Juanda showing a dragonfly to granddaughter Molly, Gainesville, Florida, June 1990

members collected it during the 1995 National DSA meeting. No such luck this time! We didn't even see *Argia tezpi* at Roy Beckemeyer's Cottonwood Draw site which was practically dried up and pretty well trampled by cattle and javelinas, but *Argia extranea* was common here.

Continuing eastward on the Geronimo Trail, we ascended the ocotillo covered hills and crossed the state line into Hidalgo County, New Mexico. Around the Black Dam area, we got new county records for *Argia lugens*, *A. moesta*, and *Libellula pulchella*. *Sympetrum corruptum* and *Enallagma praevarum* were common in the washed-out streams and a few *Archilestes grandis* gave Bob's photographic patience a work out. I saw several colorful endemic dytiscids called the Sunburst Diving Beetle (*Thermonectes marmoratus*) swimming away from hungry *Pantala* larvae! On the way back to town, we stopped at a nearby cattle pond off Guadalupe Canyon Road. There, we got new Cochise County records for *Pantala hymenaea* and the spooky *Tramea onusta*. *Lestes alacer* was very common among the dried shrubs. At one little puddle, I saw several diminutive red-spotted toads (*Bufo punctatus*), a species I had never seen before.

The next day, we went to the Slaughter Ranch and the San Bernardino NWR. Bob and Steve looked for *Argia "carlcooki?"* and *Enallagma semicirculare* which Bob had photographed here last year. Unfortunately, as Bob put it, "We didn't even see a vapor trail of either species!" A conversation with Nina King, the Refuge Manager, confirmed that odonate and other invertebrate populations were down this summer. Meanwhile, Bill and I had better luck at the famous Slaughter Ranch Pond. Swarming dragonflies and damselflies were teeming over the water! Such highlight species as *Aeshna dugesi*, *Coryphaeschna luteipennis florida*, and *Perithemis domitia* were flying among the hordes of *Aeshna multicolor*, *Libellula saturata*, *Pachydiplax longipennis*, and *Enallagma praevarum*! I got about 25 species altogether but I would have given them all up for one specimen of a giant *Aphylla*-like gomphid that cruised along the west bank of the pond that morning. I saw it 3 different times but it never landed. The club was red and yellow striped. I don't know what it could be. Does anyone have any ideas? I remember getting a glimpse of one in 1995 but I wasn't sure. Since nobody caught one, a return trip is in order for next year!

The next day, at Bob's urging, we visited the famous Muleshoe Ranch located west of Willcox. This beautiful place is mostly managed by the Nature Conservancy. Having contacted the ranch well in advance, we were able to collect at Hot Springs Canyon, one of six perennial flowing streams there in the Galiuro Mountains.

This place is a desert Paradise with a high diversity of odonates, butterflies, birds, reptiles and other native wildlife! We collected close to 30 species of odonates, including such highlights as: *Apanisagrion lais*, *Argia oenea*, *A. pima*, *A. tarascana*, *Palaemnema domina*, and *Erpetogomphus natrix*.

The slender, green-eyed *Apanisagrion lais* was in shady, seepage trickles and they behaved much like the tropical *Aeolagrion* or *Acanthagrion*. The red-eyed, metallic *Argia oenea* and *Erpetogomphus natrix* were very common at gravel bars and riffles. *Argia pima*, *A. tarascana*, and *Palaemnema domina* inhabited the tangled, streamside root masses of the giant cottonwoods. It was not uncommon to see 3 or 4 *A. tarascana* and *A. pima* perching on the sunny roots. After a brief noontime shower, a few *P. domina* appeared in the openings at the bases of the trees. While *Argia oenea* and *A. tarascana* were common, *A. pima* and *P. domina* were not. Hopefully, sizeable populations will be found on the other five nearby streams.

Wildlife was in abundance there at Muleshoe. A herd of javelina greeted us as we stepped down in the first dry streambed. I don't know who was more scared, us or them! We saw several pronghorn antelope on the beautiful desert prairie outside the ranch entrance; truly magnificent animals and the highlight of the trip for me! Rattlesnakes are common here and I did not know that there are at least 4 species present! Butterflies swarmed at the flowering shrubs and birds were common and included: Roadrunner, Zone-tailed (a good one to see), Red-tailed, and Gray hawks (another good one), Black Phoebe, Wilson's and Nashville warblers, both Western and Summer tanagers, and Lesser Goldfinch.

The last couple of days were spent at Tucson. On our first day we ascended the beautiful Catalina Highway into the Santa Catalina Mountains, entering through a dramatic and alien forest of spectacular giant and ancient Saguaro cactus. At Molino Prison Campground, we had to mosey through a Hell's Angels biker campout. Harley-Davidson and Kawasaki motorcycles plus their

bearded, imposing owners were everywhere! Biker women were particularly interested in what we were doing with our nets and cameras!

There was a lot more water present than last year, and *Archilestes grandis*, *Argia munda*, and *A. sabino* were common. Bob and Steve got excellent photos of *A. munda*, a desert species rarely seen in the United States.

At noon, we went drove further up the Santa Catalina Mountains to Summerhaven at about 9,000 feet. At Marshall Gulch Creek, we got several giant, pale *Oplonaeschna armata*. All the specimens were young but in good shape and this may be due to the cold water delaying their summertime emergence.

After the next morning's breakfast, Bob left us to visit friends in Silver City as he drove back to Houston. The rest of us worked Lower Sabino Canyon because Upper Sabino Canyon and several of the tram bridges were wiped out by torrential July storm flooding! The storm filled all the deep *Argia sabino* pools with sand and knocked down all the riparian trees and vegetation! It looked like a defoliated war zone! Steve and I saw one, maybe two, *A. sabino* but my dependable *A. pima* site was totally gone! It will be several years before Sabino Canyon is back to normal. Recolonization by *A. sabino* is possible since we'd seen several adults the previous day upstream at Soldier Canyon near the Molino Prison Campground. However, I think *A. pima* may be extirpated from Sabino Canyon.

Lower Sabino Canyon was a different story! Swarming dragonflies were everywhere! All that flooding created several flowing streams, deep pools, and desert marshes. *Erythrodiplox connata* [Dennis Paulson is currently preparing an article proposing a return to the Calvert name *basifusca* for this species. Ed.], *Anax walsinghami*, *Argia pallens*, and various libellulids were very common. We even went back the next day to catch the elusive *Dythemis nigrescens* (?) that was giving Steve and me fits before getting away!

All good things must come to an end. The next day, the Rat Patrol parted company and headed home. It was a blast and we are ready to do it again! I had a couple of days left, so I headed north to Pine Flat Campground in Oak Creek Canyon near Flagstaff, Arizona.

Although most of the deep pools were filled with gravel and sand from past storms, I did find two

being visited by the aptly and beautifully named *Aeshna persephone*, my favorite aeshnid! I was able to observe females ovipositing in mosses on boulders in the middle of the creek. The wary males began patrolling in the afternoon around 4:00pm but I did manage, after many wild swings, to successfully net a few of these magnificent dragons! Aside from a few *Archilestes grandis* and *Argia tonto*, there were no other odonates on the creek.

All in all, it was a great trip! Everyone had a blast and we are looking forward to exploring new desert country next year! Adios, muchachos!

=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=

NEW RECORDS OF ODONATA FOR SOME PROVINCES OF THE DOMINICAN REPUBLIC

José M. Ramos Hernandez, Apartado Postal 2204, Sancti-Spiritus, CUBA, CP 60100
[Translated by Nick Donnelly]

Between February and March 1999, Dr. Luis F. de Armas of the Instituto de Ecología y Sistemática of La Habana made a trip to the Dominican Republic. In addition to collecting scorpions, he also spent part of his time collecting butterflies, spiders, insects, etc. In this last group the odonates were well represented, with 77 specimens of 15 species, whose list we present, including new provincial records for the Dominican Republic.

RESULTS – Zygoptera, Coenagrionidae

Enallagma coecum (Hagen): New record for Monte Plata Province; also collected in the Distrito Nacional.

Telebasis vulnerata Hagen: New records for Monte Plata and Monseñor Nouel Provinces.

Megapodagrionidae

Hypolestes trinitatis Gundlach: Collected in the Distrito Nacional.

Anisoptera, Aeshnidae

Triacanthagyna trifida (Rambur): Collected in La Vega Province.

Libellulidae

Dythemis rufinervis (Burmeister): New record for Monte Plata Province; also collected in the Distrito Nacional, and Santiago.

Erythrodiplax umbrata (Linn.): New record for Puerto Plata Province; also collected in Monseñor Nouel, Pedernales and Distrito Nacional.

Erythemis vesiculosa (Fabr.): New record for Monte Plata Province.

Macrothemis celeno (Selys): New record for Monte Plata and Puerto Plata Provinces; also collected in Monseñor Nouel and Distrito Nacional.

Micrathyria aequalis (Hagen): New records from Pedernales and Monte Plata Provinces.

Micrathyria dissocians Calvert: New record for Monte Plata Province.

Micrathyria hagenii Kirby: New record for Monte Plata Province.

Orthemis [ferruginea] (Fabr.): Collected in Puerto Plata Province. [Note: Antillean *Orthemis* assigned to this species represent an undescribed species currently being studied. Ed.]

Pantala flavescens (Fabr.): Collected in Puerto Plata Province.

Perithemis domitia (Drury): Collected in the Distrito Nacional.

Scapanea frontalis (Burmeister): Collected in La Vega Province and the Distrito Nacional.

Acknowledgments: I am sincerely grateful to Dr. Luis F. de Armas of the Instituto de Ecología y Sistemática for having furnished me with the odonate material collected by himself in the Dominican Republic for this study; to Dr. Oliver S. Flint, Jr. of the National Museum of Natural History in Washington for having sent to Dr. de Armas part of the bibliography used in this study; to Dr. Thomas Donnelly, Dragonfly Society of the Americas, New York, and to Bill Mauffray, of the International Odonata Research Institute, Gainesville, Florida, for having sent part of the literature utilized here.

Bibliography

Alayo, D., P., 1968. Las Libélulas de Cuba (Insecta: Odonata). *Torreia*, n.s. 2: 1-102 (Part I: text), 3: 1-54 (Part 2, plates)

Daigle, J.J., 1993. Checklist of the Odonata of the Dominican Republic by province. *Bulletin of American Odonatology* 1(4): 65-69.

Daigle, J.J. New records of Odonata for the Dominican Republic. ARGIA

Westfall, M.J., Jr., and M.L. May., 1996. *Damselflies of North America*. Gainesville, Scientific Publishers, 650 pp.

ERYTHEMIS VESICULOSA (FABRICIUS), GREAT PONDHAWK, NEW FOR ALABAMA

R. Stephen Krotzer, 6010 Woodvale Drive, Helena, AL 35080, smjkrotzer@aol.com

During 1999, I was contracted by the Alabama Department of Conservation and Natural Resources Natural Heritage Section to conduct an odonate survey of the Grand Bay Savanna Nature Preserve, a part of the state's Forever Wild Program's land holdings in Mobile County, Alabama. On 14 August between 1000 and 1230 hours, while conducting this survey, I collected one mature male *Erythemis vesiculosa* and sighted at least one other male. These individuals were located along a transmission line right-of-way in the immediate vicinity of a number of small depressional pools and wet ditches, which appeared to be permanent or semi-permanent in nature. Although the site was visited at least monthly between April and September, no additional individuals of the Great Pondhawk were encountered.

Erythemis vesiculosa is a fairly common tropical species that has been reported from a number of states. Dunkle (1989) lists it as common in south Florida, occurring along the Gulf Coast north to Fort Myers; he also includes Oklahoma and Arizona as historical collection localities for the species. The Great Pondhawk has also been collected in Kansas, New Mexico, and Texas (Bick & Mauffray, 1997-1999). This new record for *E. vesiculosa*, then, appears to be the first documented occurrence of the species in the area along the Gulf Coast between the Texas/Louisiana border and Fort Myers, Florida. Whether these individuals are simply vagrants, or whether the species is slowly colonizing suitable habitat along the Gulf Coast, is anybody's guess at this point.

The habitat and conditions where the Great Pondhawk specimen was found are not what I normally associate with "primo" dragonfly collecting – low habitat diversity with mostly shallow, hot water; daytime temperatures over 100 degrees with nearly 100% humidity; odonate abundance and diversity low; deerfly and mosquito abundance extraordinarily high!! If I hadn't been working under contract, I would not been out there collecting at high noon in August, and I almost certainly would not have gotten this noteworthy record. For that reason, I'd like to extend a special thanks to Greg Lein, Natural Heritage Section Chief; the funding he provided helped ease the sting of all those insect bites (not to mention the speeding ticket)!!

REFERENCES

Bick, G.H. and B. Mauffray. 1999. Distribution Summary of North American Anisoptera (B. Mauffray and L. Beckenbach eds., 1997). Available on the Internet at: <http://www.afn.org/~iori/nalist.html>

Dunkle, S.W. 1989. Dragonflies of the Florida Peninsula, Bermuda and the Bahamas. Scientific Publishers, Gainesville, Florida. 155 pp.

==||==||==||==||==||==||==||==||==||==||

MOLECULAR PHYLOGENETIC ANALYSES OF THE ODONATE GENERA *LIBELLULA*, *LADONA* AND *PLATHEMIS*

THOMAS ARTISS

The odonate genus *Libellula* Linnæus 1758, has been the focus of considerable taxonomic controversy. The central issue of debate revolves around whether *Plathemis* Hagen, 1861 and *Ladona* Needham, 1897 should be accorded generic status, be considered subgenera within *Libellula* s.l., or be synonymized with *Libellula* s.s. These groups were assigned full generic rank by Needham (1897), and this interpretation has been widely adopted (Needham and Westfall 1955; Carle 1978; May 1992; Schmidt 1987; Westfall and Tennenssen 1996). Several other authors, however, have suggested that these two groups be considered subgenera within *Libellula* s.l. (Kennedy 1922a,b; Borror 1945; Bennefield 1965; Walker and Corbet 1975; Allen et al. 1985; Tsuda 1986). Still others have concluded that *Ladona* and *Plathemis* are not subgenera, and have synonymized them with *Libellula* s.l. (Ris 1910; Garman 1927; Byers

1930). The situation is further complicated by the uncertain assignment of two European *Libellula* species, *L. depressa* and *L. fulva*, that have been assigned to *Plathemis* and *Ladona* respectively by some authors (Needham 1897; Needham and Westfall 1955; Schmidt 1987), included within *Libellula* s.s. by others (Tsuda 1986), or even given their own subgeneric ranks (Kennedy 1922 a,b).

One of the principle objectives of my dissertation research is to examine taxonomic questions regarding relationships among taxa within the genus *Libellula*, and to determine whether members of this genus formed a natural, or monophyletic clade (monophyletic clades are all organisms that share a unique, common ancestry). Only one other study has examined these taxa using robust phylogenetic techniques (Kambhampati and Charlton 1999). This study concluded that *Plathemis* and *Ladona* were distinct, monophyletic lineages that were basal within *Libellula* s.l.. However, this study was unable to find strong resolution within *Libellula* s.s., and did not survey several species in the genus, including *L. fulva* and *depressa*.

DNA from 26 species of *Libellula* and 4 outgroup taxa was isolated, amplified and used to collect sequence data from two mitochondrial genes (cytochrome oxidase I and 16S rRNA). Mitochondrial DNA has the distinct advantage that it is inherited only from the maternal parent and without recombination in most organisms, unlike nuclear genes that experience recombination each generation. Thus mitochondria provide a direct chain of ancestry across generations. Moreover, because the selective pressures that act on nucleotides seem to be of a different nature than those which act on morphological characters, DNA sequences provide a very valuable additional data set which can aid in resolving situations where morphological characters sets are simply unsuitable due to the nature of selection acting on them. These particular mitochondrial genes were selected because they have been widely used for invertebrate molecular systematics, and the rate of evolution of these genes is suitable for species-level phylogenetic studies.

These data were analyzed using three principle methods of phylogenetic analysis; minimum evolution, parsimony and maximum likelihood. Each of these methods are philosophically different approaches to phylogenetic reconstruction, however it is generally agreed that confidence in a tree topology is enhanced if each method independently produces trees that are generally similar. Briefly,

minimum evolution analyses evaluate numbers of changes in sequences between taxa, and fit them to a tree incorporating a model of evolution, maximum parsimony methods create trees that requires the smallest number of changes to explain observed differences in taxa, and maximum likelihood analyses produce the most probable tree under a specific model of evolution. Details of specific analyses and protocols will be available in a forthcoming manuscript.

The results of a combined data set analysis are presented in Figure 1. There are three important conclusions from this study.

(1) The genus *Orthemis* is traditionally distinguished from *Libellula* on wing venation characters. However, my results indicate that the monophyly of *Libellula* was not supported, and *Orthemis ferruginea* was found to be part of *Libellula* s.l. I employed statistical tests to determine whether this tree was significantly different from trees where *Orthemis* was constrained to be an outgroup to *Libellula* s.l. There were no significant differences in these trees. Given the indecisiveness of the molecular data on this point, I suggest that, because it is conventionally favored by morphological data, we should continue to accept the current hypothesis of *Libellula* monophyly and a corresponding outgroup position for *O. ferruginea*. However, I caution that traditional classification of these taxa may be based on potentially homoplastic (similarities not due to common ancestry) characters, and suggest that further research is needed to assess the natural delineations of taxa in these groups.

(2) The results of my study indicate that *Plathemis* and *Ladona* are distinct monophyletic lineages within *Libellula* s.l. My results indicate that *Plathemis* forms the basal sister group to the remainder of *Libellula* s.l., and that *Ladona* is the next most basal clade within the *Libellula* lineage. All three analytical approaches supported the monophyly of *Plathemis* and *Ladona*, and there was strong quantitative support (bootstrap values >90%) for these groups. Bootstrap values indicate the percent support for clades based on the randomization tests (a bootstrap value of 100 indicates that clade was supported in 100% of randomization tests). These results therefore support the original classification proposed by Needham (1897) in regard to the delineation of natural groups, with the exception that *Libellula depressa* and *Libellula fulva* would both need to be included within *Ladona* s.l. were it accorded

generic or subgeneric status. Based on my results, I propose that the separate generic or subgeneric ranks be adopted for *Plathemis* and *Ladona* within Libellulidae; a conclusion that was supported by a previous molecular phylogenetic study on these groups (Kambhampati and Charlton 1999).

(3) Phylogenetic relationships within *Libellula* s.s. generally supported the subgeneric classifications of Kennedy (1922 a,b). While Kennedy proposed that *L. angelina*, *semifasciata* and *quadrimaculata* were separate sub-genera, I found strong support for these species forming a monophyletic clade. I also found support for the subgenera *Neotetrum* (*forensis*, *pulchella*, *nodistica*), *Belonia* (*foliata*, *saturata*, *croceipennis*), and *Holotania* (*axilena*, *composita*, *jesseana*, *flavida*, *auripennis*, *luctuosa*, *cyanea*, *comanche*, *incesta*, *vibrans*). The only exception to Kennedy's classification was *L. composita* which he assigned to *Holotania*, but I found to be part of, or sister taxon to *Belonia*.

The research from this study will assist in our understanding of the diverse morphology, ecology, behavior and distribution of *Libellula* species, and future studies on the evolution of traits in this genus should be conducted with reference to phylogenetic framework. This study represents a critical step in determining the historical relationships among members of the genus, and establishes a framework for subsequent comparative studies of behavior and evolution within the group.

Literature Cited

- Allen, D., Davies, L., and Tobin, P. (1985). The dragonflies of the world: A systematic list of extant species of Odonata. Vol. 2 Anisoptera. Soc. Int. Odonatol. Rap. Comm. Suppl. 5.
- Bennefield, B.L. (1965). A taxonomic study of the subgenus *Ladona* (Odonata: Libellulidae). Univ. Kans. Sci. Bull. 45: 361-396.
- Borror, D.J. (1945). A key to the New World genera of Libellulidae. Ann. Entomol. Soc. Am. 38: 168-194.
- Byers, C.F. (1930). A contribution to the knowledge of Florida odonata. Univ. Florida Publ. Biol. Sci. 1: 1-327.
- Carle, F.L. (1978). Progress on taxonomic and nomenclatural problems of Virginia Anisoptera: a reply to Dennis R. Paulson. Selysia 8(1): 13-14.
- Garman, P. (1927). The Odonata or dragonflies of Connecticut. Guide to the Insects of Connecticut Part V. Conn. Geol. and Nat. Hist. Surv. Bull. 39: 1-331.





Figure 1. Strict consensus tree of two trees inferred from a maximum likelihood analysis based on COI and 16S data. Bootstrap values (based on 100 replicates) >50% shown above branches. Putative sub-genera assigned by Kennedy (1922 a,b) indicated to the right. Dashed line around *Belonia* indicates that *Libellula composita* was not assigned to this sub-genus by Kennedy.



Kennedy's *Libellula* (senso latu) phylogeny, 1922

- Kambhampati, S. and Charlton, R.E. (1998). Phylogenetic relationships among *Libellula*, *Ladona* and *Plathemis* (Odonata: Libellulidae) based on DNA sequence of mitochondrial 16S rRNA gene. Syst. Ent., 24: 37-49.
- Kennedy, C.H. (1922a). The morphology of the penis in the genus *Libellula* (Odonata). Ent. News 33: 33-40.
- Kennedy, C.H. (1922b). The phylogeny and the geographical distribution of the genus *Libellula* (Odonata). Ent. News 33: 65-71.
- May, M.L. (1992). Morphological and ecological differences among species of *Ladona* (Anisoptera: Libellulidae). Bull. Am. Odonat. 1(3): 51-56.
- Needham, J.G. (1897). *Libellula deplanata* of Rambur. Can. Ent. 29: 144-146.
- Needham, J.G. and Westfall, M.J. (1955). "A manual of the dragonflies of North America (Anisoptera)" University of California Press, Berkeley.
- Ris, F. (1910). Collections zoologiques du Baron Edm. de Selys Longchamps. Catalogue systematique et descriptif. Fascimile XI. Libellulinen 3 1: 245-384 + 1 pl.
- Schmidt, E. (1987). Generic reclassification of some Westpalaeartic Odonata taxa in view of their Nearctic affinities (Anisoptera: Gomphidae, Libellulidae). Adv. Odonat. 3: 135-145.
- Tsuda, S. (1986). "A Distributional List of World Odonata" Osaka.
- Walker, E.M. and J.S. Corbet (1975). "The Odonata of Canada and Alaska, Vol. 3" University of Toronto Press, Toronto.
- Westfall, M.J., Jr. and Tennessen, K.J. (1996). Odonata. In "An Introduction to the Aquatic Insects of North America, 3rd Ed." (R.W. Merritt and K. W. Cummins, Eds.), pp. 164-211. Kendall/Hunt Publishing Company, Dubuque, Iowa.

=| =| =| =| =| =| =| =| =| =| =| =| =| =| =| =|

HISTORY OF AMERICAN ODONATA: CLARENCE KENNEDY (1879-1952)

Nick Donnelly

Clarence H. Kennedy was a remarkable and productive student of the Odonata from the turn of the century to the late 40's. He was born in Rockport, Indiana, in 1879. Little has been recorded of his boyhood years, but he met E.B. Williamson around the turn of the century and the older Williamson (by two years) undoubtedly had a great influence on Kennedy. Kennedy graduated from Indiana State University in 1903, having

already written two papers: one on the use of the mesostigmal lamina for identifying females of the difficult damselfly genus *Argia*, and the second on the Odonata of Winona Lake, Indiana (where he evidently taught at a bible camp). In these papers he makes it obvious that he was heavily influenced by Williamson, who had visited him at Winona Lake. He worked briefly during this period for the U.S. Fisheries Commission, where he illustrated several papers on eel taxonomy.

After receiving a masters degree from Indiana State in 1903, his activities were not recorded. Evidently he fell ill during that period, and during his illness he met a Dr. Davis, a combination ophthalmologist, osteopath, and homeopath. In 1908, he went to Texas, where Davis was located, and then followed him to Oregon. Kennedy was seriously handicapped by illness during his earlier years, which probably explained his somewhat disjointed early life and also the completion of his doctorate at the age of forty. It seems that illness brought him to the venue of his greatest odonatological success: the Pacific Northwest.

Kennedy was an all-round naturalist, and his bibliography contains numerous early papers on birds ("Some robin's and mourning dove's nests in the lower Yakima valley. . .", "Notes on the fruit-eating habits of the sage thrasher. . .")

Between 1909 and 1914 he collected dragonflies in Oregon and Washington. During part of this period he was employed by the U.S. Bureau of Fisheries. In a three-year period (1911 to 1914) he burst forth with a trilogy of papers on the western Odonata fauna that established him as one of the foremost Odonatists in America.

On 11 and 12 July 1911, while engaged in a Mussel survey at Bumping Lake in Yakima County, Washington, he collected four species of Odonata, including a long series of the Emerald, *Somatochlora semicircularis*. Because this species had been poorly known, he wrote a paper ("The Odonata of Bumping Lake", 1913) redescribing the species, and producing the first of what came to be known as some of the finest Odonata illustrations ever produced. The paper contains no less than 57 figures illustrating the variation he observed in his collection. The remainder of the Washington and Oregon work was published as "Notes on the life history and ecology of the dragonflies of Washington and Oregon (1915)". In this profusely illustrated paper (200 figures) he described *Argia emma* (named for his mother) and figured and

discussed adults and larvae of many hitherto little known northwestern species.

During the summer of 1914 he made several trips through central California and Nevada. Kennedy paid for his trips with a railroad pass provided by the director of the California Academy of Sciences, money earned illustrating birds, insects, and fish, and a subvention from his old friend Williamson, "who later received his pay in a series of the specimens collected." The trip, as we can partially reconstruct it from the serial account, must have been one of the all-time Odonata adventures in the United States. Fortunately the editor of the Proceedings of the National Museum did not red-pencil his frequent comments about the habits of the insects. I quoted his remarks about Pyramid Lake, Nevada, in a previous ARGIA. Another comment concerns *Aeshna multicolor*, observed in Sacramento: "This species was observed catching insects on the market street of the city at twilight. They flew among the wagons and buggies entirely indifferent to the numerous passers-by. This habit of familiarity with man's haunts is very noticeable in *multicolor*. It is the most domestic of all the western Odonata."

The paper that resulted from this trip ("Notes on the life history and ecology of the dragonflies of central California and Nevada.", 1917) is still regarded as a landmark paper in North American Odonata. In this magnificently illustrated paper (more than 400 of his now famous figures), he named three new species and erected several new genera. His descriptions and illustrations for several little-known species are still consulted by workers in the far West. He also described many larvae, especially of gomphines.

His manuscript for the earlier Washington and Oregon work was with him when he went to Stanford University in the Fall of 1915 to apply for graduate study. Professor Vernon Kellogg, the Chair of the department, was so impressed by his manuscript that he awarded a masters degree on the spot to Kennedy and advised to keep on studying Odonata. He described several western *Argia* species in later papers and wrote a fine summary (well illustrated, as always) of the Odonata of Kansas (1917).

He later went to Cornell, where he obtained his doctorate in 1919 with a dissertation on the phylogeny of the Zygoptera. In this paper he made extensive use of the penile organ to help deduce phylogenetic relationships among the damselflies.

Phylogenetic deduction consumed a large part of his later life, and he later used studies of the penes to deduce relationships (and putative phylogenies) of the Neotropical damselfly genus *Acanthagrion*, and of the North American genus *Libellula*. Because this issue of ARGIA contains a more modern study of the same group using DNA (Artiss, this issue), I cannot resist including a reproduction of Kennedy's phylogenetic tree to compare with Artiss' results.

Kennedy was intensely interested in fossil Odonata and used paleontological examples to support his phylogenetic ideas. He firmly believed that the beginnings of phylogenetic stems were the most generalized forms, and also that simpler forms were more primitive than evolved. He was not alone in this line of belief, but the school of phylogeny that he represented has largely been supplanted by more modern approaches. However, it is instructive to see how many of his groupings still stand up well to modern analysis, and I wonder if a half century from now people will be referring disparagingly to the views of phylogeny that are being pursued now.


Kennedy joined the staff of the Department of Entomology of Ohio State University in 1919, becoming a full professor in 1933. His physical handicaps forced him to occupy a half-time position initially, but his health improved markedly later in life. He retired in 1949 and died in 1952.

Throughout his life Kennedy was noted as one of the finest illustrators of Odonata that has appeared. His first paper (1903) on *Argia mesostigmal laminae* has simple but elegant illustrations (nicely drawn but badly reproduced), and his 1913 Bumping Lake paper has no fewer than 57 excellent figures of one species - *Somatochlora semicircularis*. During and after World War II he produced a series of papers on South American material (mainly damselflies) that had been collected by a professional collector. The excellence of the illustrations in these papers probably added considerably to his reputation as a taxonomist.


Throughout his career Kennedy was noted for his painstaking work, and he became one of the giants of American Odonata study. His reputation in the field of entomology was considerable. He was managing editor of the Annals of the Entomological Society of America for 16 years and was president of the Society in 1935. It is a little known and somewhat sad fact that towards the end of his



The younger Kennedy – Indiana Jones or what?



One of Kennedy's famous book plates. Perhaps he didn't sleep comfortably. . .



Another book plate. For those not familiar with tropical collecting, *Triacanthagya* is one of the famous "night fliers": darners which appear at dusk and fly erratically, testing the skill of collectors.



The older Kennedy

In view of the development planned for Cypress Provincial Park, the future of this, the only known breeding site for *T. hageni* in Canada, is uncertain.

=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=

MINUTES, DSA MEETING, PAUL SMITHS, NEW YORK, 8 -10 July 1999

Sid Dunkle, DSA Secretary

Attendees included: John Abbott; Jeff Ballard; Bob Barber; Roy and Pat Beckemeyer; Ethan Bright; Everett Cashatt; Carolyn Cass; Duncan Cuyler; Dave Czaplak; Nick and Ailsa Donnelly; Sid Dunkle; Pam Guy; Bob and Jacob Harding; George and Phoebe Harp; Gord Hutchings; Dan and Karol Lynn Johnson; Jim Johnson; Gerhard Jurzitza; Ellis Lauder milk; Larry Master; Bill Mauffray; Mike and Leslie May; Don Miller; Blair Nikula; Paul Novak; Mark, Adrienne, and Marjorie O'Brien; Dennis Paulson; Eric Pilgrim; Werner and Misa Piper; Teresa Prather; Martha Reinhardt; Clark Shiffer; Fred and Peggy Sibley; Joe Smentowski; Ken Soltesz; Dan Spada; Jeremiah Trimble; Jeanette Tromblay; Jan Trybula; Tim Vogt; and Hal White.

President Michael May called the business meeting to order and read a statement from past president Rosser Garrison, who was unable to attend. Dr. May mentioned three things the DSA can do to advance odonatology: (1) Collate data on Dragonfly Migration. (2) Archive data on the Habitat Requirements of individual Odonata species; volunteers are needed to coordinate this, and the data will be placed on the IORI Website by Bill Mauffray. (3) Define the ranges of North American Odonata via Dot Maps, which is already being done by Nick Donnelly.

Some Bylaws of the DSA need to be changed, which will be done first by the DSA Council, then voted on by the membership, probably at the next meeting. The minutes of the 1998 DSA meeting were approved. The new President-Elect is Dennis Paulson. Dennis discussed possible meeting sites for the next DSA Meeting, and it was decided that the meeting will be in Vancouver, British Columbia, probably in July 2000, organized by Gordon Hutchings. It is likely that the Okanagon River Valley can also be visited as part of that meeting. Suggestions are needed for the 2001 and 2002 meetings, which can be discussed on ODONATA, Dennis Paulson's list-serve.

Nick Donnelly, editor of ARGIA and Bulletin of American Odonatology, stated that ARGIA can

use colored photos for covers, while more manuscripts are needed for BAO. The state surveys presented as issues of BAO have gone well, and the North American Odonata Dot Maps will probably also be published in BAO. Treasurer Jerrell Daigle said that DSA began 1999 with \$11190.06 in its treasury, and currently has \$13412.55. The only current expenses for DSA are the production and mailing of ARGIA (300 subscribers) and BAO (190 subscribers). After estimated expenses of \$8000, DSA is expected to end 1999 with about \$7000. Dan Johnson announced that recovery plans for the federally endangered species Hine's Emerald (*Somatochlora hineana*) are available for review by those who will return comments, pro or con, to him.

Evening presentations included: Nick Donnelly, tropical odonates from Brazil and SE Asia; Gerhard Jurzitza, *Sympetrum*s in flight; Tim Vogt (with Tim Cashatt and James Purdue), Phylogeny of *Somatochlora*; Hal White, catching *Epiaeschna*; and Dennis Paulson, Odonata of Australia.

=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=

1999 DSA FINANCIAL REPORT

Jerrell J. Daigle

At the request of Nick Donnelly, I have prepared a short summary of the ARGIA and BAO financial account for 1999. Dues for both journals are combined together in one savings account at the SunTrust Bank in Tallahassee, Florida. We began the 1999 year with a 1998 balance forward amount of \$11,190.06.

Presently, our current account is \$12,138.96. Our expenses were for ARGIA 10(4), 11(1), 11(2), and BAO 5(4). Annual expenses were estimated at \$8,000.00 for all ARGIA and BAO mailings. After all expenses and without any incoming late dues for 1999, our projected surplus by the end of 1999 is estimated at \$7,000.

A brief current financial report was presented at the July DSA meeting in Paul Smiths, New York and a few copies of the report were distributed to the media.

Also, ARGIA and BAO subscriptions have been increasing dramatically. According to Ailsa Donnelly, we have about 315 ARGIA and 105 BAO subscribers. Like soaring *Aeshnas*, we can go only higher in the next millennium!

=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=|=

CORRECTION - The Tennessee field meeting account regrettably omitted the name of one of the participants - Jerrell Daigle's father, Jerry, who helped in the finding of several of the interesting species, including the state record *Ischnura kellicotti*.

NEW OLD PUBLICATION ON SOUTH FLORIDA ANISOPTERA.

Dennis Paulson has just published his PhD dissertation from 1966 on The Dragonflies (Anisoptera) of South Florida. This 139-page publication (Slater Museum of Natural History Occasional Paper No. 57) contains much information on the status of South Florida dragonflies and their natural history. To receive a copy, send a check for \$12.00 made out to the University of Puget Sound to Dennis Paulson, Slater Museum of Natural History, University of Puget Sound, Tacoma, WA 98416.

TRAMEA

Some contributions from **Roy Beckemeyer**:

Robert Larsen, a cooperative research biologist at the Bitter Lake NWR near Roswell, New Mexico, has a site on the web that comprises an Odonata checklist for New Mexico. It is based on Mary Alice Evans' 1995 published checklist, supplemented by records from a little known and previously neglected work by Harold B. Freshley on the Upper Pecos Watershed (1945 University of New Mexico Thesis), and by records supplied by Mr. Larsen and by a number of members of this society. This very useful set of pages contain lists of counties for each species, and a map for keying out county locations, and is available at:
<http://www.rt66.com/~kjherman/odonata/NMdrngfly.html>

Cathy Biggs, who is doing much hard work to document the dragonfly distribution in California, has a web page on which she describes a spiral bound color booklet called "Common Dragonflies of California" which she is planning to eventually publish located at:
<http://www.sonic.net/~bigsnest/Pond/Lists/COVER.html>

On the international scene, check out:

<http://home9.swipnet.se/~w-90582/dragonfly/dragonfly.html>

This is Martin Petersen's Swedish Dragonfly page. Quite remarkable and out of the ordinary are: Dragonfly games, Multilingual checklists, and a dragonfly calendar.

"Dragonflies and damselflies of Kuala Lumpur",

<http://www.geocities.com/RainForest/Vines/8983/dragonflies/dragonfl.html>

by Chin Fah Shin, has a number of color photos of colorful odonates from the Orient.

And, while we are thinking oriental, why not try:

<http://www.dfw.net/~jazzman/knotter/dragon.htm>

to see directions on how to tie a Chinese ornamental dragonfly knot? Directions are illustrated step-by-step, and include a photograph of examples tied by author Jack Keene and by Tzu-ni Yang and Ming Jyh-Chen of Taiwan. These little dragonflies could be neat additions to holiday gift packages.

BACK ISSUES OF ARGIA AND THE BULLETIN OF AMERICAN ODONATOLOGY

The editor is able to provide back issues of ARGIA. Please contact T. Donnelly, 2091 Partridge Lane, Binghamton NY 13903. The present price schedule takes into account the different costs of duplication of each number of ARGIA. In the event that an issue becomes exhausted, then xerox copies will be sent. Prices are \$2.00 per issue; these do not include postage; see below.

Back Issues of the BULLETIN OF AMERICA ODONATOLOGY can be furnished at the prices given below. Prices do not include postage; see below.

1(1) The Odonata of New York, Thomas W. Donnelly p. 1-28	\$3.00
1(2) Distribution of Dragonflies and Damselflies in Florida, Sidney W. Dunkle p. 29-50	\$2.50
1(3) Morphological and ecological differences among species of <i>Ladona</i> , Michael L. May p. 51-56	\$1.50
Comportamiento reproductivo y policromatismo en <i>Ischnura denticollis</i> Burmeister, Alejandro Córdoba Aguilar [with English summary] p. 57-64	
1(4) A checklist of the Odonata of the Dominican Republic by Province, Jerrell James Daigle p. 65-69	\$1.50
Odonata de la Sierra de Huauchinango, Puebla, Mexico [with English summary], José A. Gómez Anaya and Rodolfo Novelo Gutiérrez p. 71-73	
2(1) La Nayade de <i>Archilestes latialatus</i> Donnelly, 1981 [with English summary], R. Novelo-Gutiérrez p. 1-7	\$1.50
Descripción e Historia Natural de las Larvas de Odonatos de Costa Rica. III <i>Gynacantha tibiata</i> (Karsch 1891) [with English summary], Alonso Ramírez p. 9-14	
2(2) Description of the Nymph of <i>Epietheca (Tetragoneuria) spinosa</i> (Hagen, K. J. Tennessen p. 15-19	\$1.50
The Larva and Adult Male of <i>Somatochlora georgiana</i> Walker, Jerrell J. Daigle p. 21-26	
2(3) <i>Macromia illinotensis</i> and <i>georgina</i> : a Study of heir Variation and Apparent Subspecific Relationship, T.W. Donnelly, K.J. Tennessen p. 27-61	\$3.00
2(4) The Subgenus <i>Tetragoneuria (Anisoptera: Corduliidae: Epietheca)</i> in New Jersey, Michael L. May p. 63-74	\$1.50
3(1) The Odonata of Ohio - a Preliminary Report, Robert C. Glotzhober p. 1-30	\$3.00
3(2) Four Decades of Stability and Change in the Odonata Population at Ten Acre Pond in Central Pennsylvania, Clark N. Shiffer & Harold B. White p. 31-40	\$1.50
Descripción e Historia Natural de las Larvas de Odonatos de Costa Rica. IV. <i>Mecistogaster ornata</i> (Rambur, 1842) [with English summary], Alonso Ramírez p. 43-47	
3(3) The Distribution of Odonata in Alabama, Kenneth J. Tennessen, James D. Harper, R. Stephen Krotzer, p. 49-74	\$3.00
3(4) Distribution Records of the Odonata of Montana, Kelly B. Miller and Daniel L. Gustafson, p. 75 - 88	\$1.50
4(1) An Annotated List of the Odonata of New Jersey, With an Appendix on Nomenclature in the Genus <i>Gomphus</i> , Michael L. May & Frank L. Carle p. 1 - 35	\$3.00
4(2) The Odonata of Patuxent Wildlife Research Center and Vicinity, Richard L. Orr p. 37 - 67	\$3.00
4(3) The Status of <i>Lestes apollinaris</i> Navás and <i>L. henschawi</i> Calvert, Thomas W. Donnelly p. 69-74	\$1.50
4(4) The Dragonflies of Washington, Dennis R. Paulson p. 75-90	\$1.50
5(1) The Dragonflies and Damselflies (Odonata) of Louisiana, Bill Mauffray p. 1-26	\$3.00
5(2) The Odonata of the Cayman Islands: a Review, R.R. Askew, R. Prosser, and P.S. Corbet p. 27-32	\$1.50
TAXONOMIC AND Population Studies of British Columbia <i>Aeshna</i> species, G. Peters p. 33-42	
5(3) Adapting the Townes Malaise Trap for Collecting Live Odonata, Robert C. Glotzhober & Dan Riggs, p. 43-48	\$1.50
<i>Archilestes grandis</i> (Great Spreadwing) in Central New Jersey, with Notes on Water Quality, David P. Moskowitz and David M. Bell, p. 49-54	
Variation in Head Spines in Female <i>Ophiogomphus</i> , with a Possible Example of Reproductive Character Displacement (Anisoptera: Gomphidae), Dennis R. Paulson, p. 55-58	
5(4) The Odonata fauna of Connecticut, David L. Wagner and Michael C. Thomas, p. 59-85	\$3.00

* subscription

Mailing and Handling Costs:

	SURFACE		AIR MAIL	
	1st number	each additional	1st number	each additional
United States	\$1.25	\$1.00	---	---
Canada, Mexico	\$1.25	\$1.00	\$1.50	\$1.25
Western Hemisphere	\$1.50	\$1.25	\$2.00	\$1.50
Europe, Asia, etc	\$1.50	\$1.25	\$3.00	\$2.50

ARGIA

Vol. 11, No. 4, 31 December 1999

Binghamton, New York

IN THIS ISSUE

VANCOUVER ISLAND DSA ANNUAL MEETING (WITH POSSIBLE OKANAGAN SIDE TRIP)	Gordon Hutchings	1
CARL COOK RECEIVES KENTUCKY AWARD	Ellis Laudermilk	2
JUANDA BICK: 1919-1999		2
ARIZONA ADVENTURES!	Jerrell J. Daigle, Bob Behrstock, Steve Krotzer, and Bill Mauffray	4
NEW RECORDS OF ODONATA FOR SOME PROVINCES OF THE DOMINICAN REPUBLIC	José M. Ramos Hernandez	6
<i>ERYTHEMIS VESICULOSA</i> (FABRICIUS), GREAT PONDHAWK, NEW FOR ALABAMA	R. Stephen Krotzer	7
MOLECULAR PHYLOGENETIC ANALYSES OF THE ODONATE GENERA <i>LIBELLULA</i> , <i>LADONA</i> AND <i>PLATHEMIS</i>	Thomas Artiss	8
HISTORY OF AMERICAN ODONATA: CLARENCE KENNEDY (1879-1952)	Nick Donnelly	12
WHAT OTHERS THINK OF US. . .		15
DRAGONFLIES IN THE NEWS		15
REQUEST FOR RECORDS FOR ODONATA IN BC	Rex D. Kenner	15
FIRST CANADIAN BREEDING RECORD FOR <i>TANYPTERYX HAGENI</i>	Rex D. Kenner	15
MINUTES, DSA MEETING, PAUL SMITHS, NEW YORK, 8 -10 July 1999	Sid Dunkle	16
1999 DSA FINANCIAL REPORT	Jerrell J. Daigle	16
CORRECTION		17
NEW OLD PUBLICATION ON SOUTH FLORIDA ANISOPTERA		17
TRAMEA	Roy Beckemeyer	17

1999 MEMBERSHIP LIST WITH E-MAIL ADDRESSES DISTRIBUTED SEPARATELY WITH THIS ISSUE