


# BULLETIN OF AMERICAN ODONATA=I=OLOGY



Volume 8 Number 1

5 Apr 2004

DISTRIBUTION OF NORTH AMERICAN ODONATA. PART II:  
MACROMIIDAE, CORDULIIDAE AND LIBELLULIDAE

Thomas W. Donnelly, p. 61 - 32

# THE DRAGONFLY SOCIETY OF THE AMERICAS

Business address: c/o T. Donnelly, 2091 Partridge Lane, Binghamton NY 13903

## EXECUTIVE COUNCIL 2003 - 2005

President	R. Beckemeyer	Wichita KS
President Elect	S. Krotzer	Centreville AL
Immediate past President	D. Paulson	Seattle WA
Vice President, Canada	R. Cannings	Victoria, BC
Vice President, Latin America	R. Novelo G.	Jalapa, Veracruz
Secretary	S. Dunkle	Plano TX
Treasurer	J. Daigle	Tallahassee FL
Editor	T. Donnelly	Binghamton NY
Regular member	J. Abbott	Austin TX
Regular member	S. Valley	Albany OR
Regular member	S. Hummel	Lake View IA

## JOURNALS PUBLISHED BY THE SOCIETY

**ARGIA**, the quarterly news journal of the **DSA**, is devoted to non-technical papers and news items relating to nearly every aspect of the study of Odonata and the people who are interested in them. The editor especially welcomes reports of studies in progress, news of forthcoming meetings, commentaries on species, habitat conservation, noteworthy occurrences, personal news items, accounts of meetings and collecting trips, and reviews of technical and non-technical publications. Articles for publication in **ARGIA** should preferably be submitted as hard copy and (if over 500 words) also on floppy disk (3.5" or 5.25"). The editor prefers MS DOS based files, preferably written in WORD, WORD for WINDOWS, WordPerfect, or WordStar. Macintosh WORD disks can be handled. All files should be submitted **unformatted and without paragraph indents**. Each submission should be accompanied by a text (=ASCII) file. Other languages should be submitted only as text (=ASCII) files. Line drawings are acceptable as illustrations.

T. Donnelly (address below) is the interim editor of **ARGIA**.

**BULLETIN OF AMERICAN ODONATOLOGY** is devoted to studies of Odonata of the New World. This journal considers a wide range of topics for publication, including faunal synopses, behavioral studies, ecological studies, etc. The **BAO** publishes taxonomic studies but will not consider the publication of new names at any taxonomic level. Enquiries and submission of manuscripts should be made to **BAO** editor T. Donnelly, 2091 Partridge Lane, Binghamton NY 13903. Final submissions (after review) should be made on floppy disk, as above, with illustrations in final form and preferably adjusted to final size.

## MEMBERSHIP IN THE DRAGONFLY SOCIETY OF THE AMERICAS

Membership in the **DSA** is open to any person in any country. Dues for individuals in the US, Canada, or Latin America are \$15 for regular membership and \$20 for institutions or contributing membership, payable annually on or before 1 March of membership year. Dues for members in the Old World are \$25.

Dues should be mailed to Jerrell Daigle, 2067 Little River Lane, TALLAHASSEE FL 32311

The **BULLETIN OF AMERICAN ODONATOLOGY** is available by a separate subscription at \$15 for members and \$18.75 for non-members and institutions.

## DISTRIBUTION OF NORTH AMERICAN ODONATA. PART II: MACROMIIDAE, CORDULIIDAE, LIBELLULIDAE<sup>1</sup>

Thomas W. Donnelly, 2091 Partridge Lane, Binghamton NY 13903, tdonnelly@binghamton.edu

**ABSTRACT:** Dot-map presentations show the distributions for species of Macromiidae, Corduliidae, and Libellulidae in North America. Additional comments, including subspecies information, are provided for several species.

**INTRODUCTION:** This is the second of a series that started with Donnelly (2004a). Introductory and explanatory comments in that number will not be repeated here.

### SUBSPECIES, AND SPECIES OF SPECIAL INTEREST:

*Macromia illinoiensis*: Donnelly and Tennessean (1994) described structural variations in the taxa *georgina* and *illinoiensis* and proposed that they had a subspecific relationship. They noted four counties where the taxa varied smoothly between the subspecific end members, and noted that in a fifth county (Macon Co. NC) there were separate populations of the two which did not so vary. May and Carle (1996) noted that "The two taxa seem distinct in structural characters in New Jersey and are sympatric in Virginia and North Carolina." We see no reason to alter our conclusion.

*Macromia X wabashensis*: I treat this as a hybrid, probably between *taeniolata* and *pacifica*. The Texas specimens might alternatively have *annulata* as one parent.

*Epitheca costalis*, *cynosura*, and *petechialis*: These taxa have caused extensive difficulties. Recent examination of numerous specimens from eastern Texas, Oklahoma, Kansas, and Colorado, have shown unspotted *petechialis* is far more widespread than previously recognized. Probably plains (western Kansas, Colorado, western Oklahoma, central and western Texas) records of "*costalis*" and "*cynosura*" are probably pale-winged *petechialis*. The only reliable method of distinguishing among these three taxa is by a close examination of the male cerci and abdominal base.

In several localities in Ohio and scattered localities elsewhere, *costalis* and *cynosura* appear to intergrade freely, although in other localities, no evidence of intergradation has been detected. While these species appear to be valid biological

species, the extent of, and the reasons for, their apparent intergradation are not yet understood.

*Epitheca semiaquea*: There is a broad gap in the distribution of this species, with no records from north-central Florida to Texas. Specimens from Massachusetts appear to lack the wing spots and have been called *cynosura*.

*Somatochlora hineae*: The records from Alabama, Ohio, and Indiana are historic.

*Williamsonia lintneri*: The record from Albany, New York is historic.

*Erythemis simplicicollis* and *collocata*: The apparent subspecific relationship between these two taxa was discussed by Donnelly (2004b).

*Leucorrhinia proxima*: The eastern and western forms are different in both appearance and habits. The eastern forms have yellow pale color and become quite pruinose. They commonly do not perch in conspicuous positions. The western forms are patterned with bright red, are far less pruinose, and perch in very exposed positions. The boundary between the two forms appears to be approximately in western Wisconsin.

*Orthemis antillean species*: This species has not been named. It flies with ferruginea around Miami, and the two can be distinguished by several differences in appearance. The antillean form ranges to Trinidad, but is not certainly specifically distinct with a co-occurring, and also unnamed, purple form in Hispaniola, Puerto Rico, and the Virgin Islands. It is premature to name any of these forms.

*Perithemis tenera*: This species appears to grade smoothly into the tropical "species" *mooma*, and several specimens from southern Texas could be referred to *mooma*. It is more realistic to regard them as one wide ranging and variable species. Some of the Mexican records shown are of "*mooma*", but Mexican records of *tenera* are numerous and widespread.

*Plathemis lydia*: Eastern and western forms, distinguished by the shape of the styli on the basal abdominal segment (Donnelly, 1991), grade

<sup>1</sup>Bulletin of American Odonatology 8(1) 1:32

together in a narrow zone in west-central Montana to western Wyoming.

*Sympetrum internum* and *rubicundulum*: Throughout most of their considerable overlapping ranges, these appear to be easily distinguishable species (Williamson, 1933). Along the Atlantic seaboard, from at least New Jersey to Cape Cod, the most abundant form appear to be *rubicundulum*, but with a hamule intermediate between this species and *internum*. Specimens of apparently valid *internum* are found sporadically in the same area.

*Sympetrum internum* and *obtrusum*: Hybrids between these two species have been found at several northeastern localities (Donnelly, 1991).

*Sympetrum semicinctum*: Specimens taken in westernmost Ontario and Wisconsin, and in Northeastern Minnesota, appear to be transitional between *semicinctum* and *occidentale*. I believe the two to have a subspecific relation, as shown. I have not further differentiated *californicum* and *fasciatum*.

#### REFERENCES CITED

- Donnelly, T.W., 1991, Current problems in *Sympetrum* and *Libellula* (a.k.a. *Plathemis*). ARGIA 3(4): 8-12

Donnelly, T.W., 2004a, Distribution of North American Odonata. Part I: Aeshnidae, Petaluridae, Gomphidae, and Cordulegastridae. Bulletin of American Odonatology 7(4): 61-90

Donnelly, T.W., 2004b, *Erythemis simplicicollis* and *collocata* – subspecies? ARGIA 15(4): 11-13

Donnelly, T.W., and Tennessee, K.J., 1994, *Macromia illinoiensis* and *georgina*: a study of their variation and apparent subspecific relationship (Odonata: Corduliidae). Bulletin of American Odonatology 2(3): 27-61

May, M.L., and Carle, F.L., 1996, An annotated list of the Odonata of New Jersey. Bulletin of American Odonatology, 4(1): 1-35

Williamson, E.B., 1933, The status of *Sympetrum assimilatum* (Uhler) and *Sympetrum decisum* (Hagen) (Odonata – Libellulinae). Occ. Papers Mus. Zoology. Univ. Michigan, no. 264, 7 p, 1 pl.

## INDEX TO MAPS

(NOTE: M refers to Macromiidae, C to Corduliidae, and L to Libellulidae)

Didymops floridensis M1  
Didymops transversa M1  
Macromia alleghaniensis M2 C1  
Macromia annulata M1  
Macromia illinoiensis M2 C1  
Macromia magnifica M1  
Macromia margarita M1  
Macromia pacifica M1  
Macromia X wabashensis M2 C1  
Macromia taeniolata M1  
Cordulia shurtleffi M2 C1  
Dorocordulia lepida M2 C1  
Dorocordulia libera M2 C1  
Epitheca (Epicordulia) princeps C3  
Epitheca (Tetragoneuria) canis C3  
Epitheca (Tetragoneuria) costalis C2  
Epitheca (Tetragoneuria) cynosura C2  
Epitheca (Tetragoneuria) petechialis C2  
Epitheca (Tetragoneuria) semiaquea C2

Epitheca (Tetragoneuria) sepia	C2
Epitheca (Tetragoneuria) spinigera	C2
Epitheca (Tetragoneuria) spinosa	C2
Epitheca (Tetragoneuria) stella	C2
Helocordulia selysi	M2 C1
Helocordulia uhleri	C4
Neurocordulia alabamensis	C3
Neurocordulia michaeli	C3
Neurocordulia molesta	C4
Neurocordulia obsoleta	C4
Neurocordulia virginiensis	C3
Neurocordulia xanthosoma	C3
Neurocordulia yamaskanensis	C4
Somatochlora albicincta	C5
Somatochlora brevicincta	C6
Somatochlora calverti	C7
Somatochlora cingulata	C5
Somatochlora elongata	C8
Somatochlora ensigera	C8
Somatochlora filosa	C8
Somatochlora forcipata	C7
Somatochlora franklinII	C7
Somatochlora georgiana	C8
Somatochlora hineana	C4


Somatochlora hudsonica	C6	Leucorrhinia patricia	L9
Somatochlora incurvata	C10 L1	Leucorrhinia proxima	L8
Somatochlora kennedyi	C7	Libellula auripennis	L10
Somatochlora linearis	C8	Libellula axilena	L10
Somatochlora margarita	C7	Libellula comanche	L10
Somatochlora minor	C9	Libellula composita	L11
Somatochlora ozarkensis	C8	Libellula croceipennis	L10
Somatochlora provocans	C9	Libellula cyanea	L10
Somatochlora sahlbergi	C6	Libellula flava	L9
Somatochlora semicircularis	C6	Libellula forensis	L8
Somatochlora septentrionalis	C6	Libellula incesta	L11
Somatochlora tenebrosa	C9	Libellula jessea	L9
Somatochlora walshii	C9	Libellula luctuosa	L12
Somatochlora whitehousei	C5	Libellula needhami	L10
Somatochlora williamsoni	C5	Libellula nodisticta	L9
Williamsonia fletcheri	C4	Libellula pulchella	L12
Williamsonia lintneri	C4	Libellula quadrimaculata	L12
Brachymesia furcata	C10 L1	Libellula saturata	L11
Brachymesia gravida	L2	Libellula semifasciata	L11
Brachymesia herbida	C10 L1	Libellula vibrans	L11
Brechmorhoga mendax	C10 L1	Macrodiplax balteata	L2
Brechmorhoga pertinax	C10 L1	Macrothemis imitans	L2
Cannaphila insularis	C10 L1	Macrothemis inacuta	C10 L1
Celithemis amanda	L5	Macrothemis inequiunguis	L2
Celithemis bertha	L5	Miathyria marcella	L5
Celithemis elisa	L7	Micrathyria aequalis	L3
Celithemis eponina	L7	Micrathyria didyma	L3
Celithemis fasciata	L7	Micrathyria hagenii	L3
Celithemis martha	L6	Nannothenemis bella	L11
Celithemis ornata	L7	Orthemis antillean species	L13
Celithemis verna	L6	Orthemis discolor	L13
Crocothemis servilia	C10 L1	Orthemis ferruginea	L13
Dythemis fugax	L2	Pachydiplax longipennis	L14
Dythemis maya	L2	Paltothemis lineatipes	L3
Dythemis nigrescens	L2	Pantala flavescens	L14
Dythemis velox	L2	Pantala hymenaea	L14
Erythemis attala	L4	Perithemis domitia	L13
Erythemis peruviana	L4	Perithemis intensa	L13
Erythemis plebeja	L5	Perithemis tenera	L13
Erythemis simplicicollis	L4	Plathemis lydia	L15
Erythemis vesiculosa	L5	Plathemis subornata	L15
Erythrodiplax basifusca	L4	Pseudoleon superbus	L5
Erythrodiplax berenice	L4	Sympetrum ambiguum	L15
Erythrodiplax funerea	L3	Sympetrum corruptum	L17
Erythrodiplax fusca	L4	Sympetrum costiferum	L16
Erythrodiplax minuscula	L3	Sympetrum danae	L16
Erythrodiplax umbrata	L5	Sympetrum illotum	L8
Idiataphe cubensis	L3	Sympetrum internum	L17
Ladona deplanata	L6	Sympetrum madidum	L15
Ladona exusta	L6	Sympetrum obtrusum	L17
Ladona julia	L7	Sympetrum pallipes	L15
Leucorrhinia borealis	L9	Sympetrum rubicundulum	L15
Leucorrhinia frigida	L7	Sympetrum semicinctum	L16
Leucorrhinia glacialis	L8	Sympetrum signiferum	L13
Leucorrhinia hudsonica	L8	Sympetrum vicinum	L18
Leucorrhinia intacta	L9	Sympetrum internum X obtrusum	L17

Tauriphila australis L4  
Tauriphila azteca L4  
Tholymis citrina L3  
Tramea abdominalis L6  
Tramea binotata L5


Tramea calverti L6  
Tramea carolina L6  
Tramea insularis L5  
Tramea lacerata L18  
Tramea onusta L18

- ◆ Adult specimen; often used for stage unspecified
- Photograph of adult
- Larva or exuviae
- ◊ Sight Record
- Record specified for state (US or Mexico) or region


Key to the symbols used in the maps.



*Didymops  
floridensis*




*Macromia  
pacifica*




*Macromia  
annulata*




*Macromia  
margarita*



*Didymops  
transversa*









*Macromia  
magnifica*



*Macromia  
taeniolata*


## MACROMIIDAE 1


*Didymops floridensis, transversa*  
*Macromia pacifica, annulata, georgina, magnifica, taeniolata*  
*margarita*




## MACROMIIDAE 2; CORDULIIDAE 1

*Macromia illinoiensis, allegheniensis, X wabashensis*  
*Cordulia shurtleffi*  
*Dorocordulia libera, lepida*  
*Helocordulia selysi*

















Many western *cynosura* records may be clear-winged *petechialis*.



Many western *costalis* records may be clear-winged *petechialis*.




## CORDULIIDAE 2    *Epitheca cynosura, costalis, petechialis, sepia, stella, spinosa*




### CORDULIIDAE 3

*Epitheca canis, spinigera, princeps*


*Neurocordulia alabamensis, michaeli, virginiensis, xanthosoma*




*Helocordulia  
uhleri*




*Neurocordulia  
molesta*




*Neurocordulia  
yamaskanensis*




*Williamsonia  
fletcheri*



*Somatochlora  
hineana*




*Williamsonia  
lintneri*




*Neurocordulia  
obsoleta*

## CORDULIIDAE 4


*Helocordulia uhleri*  
*Neurocordulia yamaskanensis, molesta, obsoleta*  
*Williamsonia fletcheri, lintneri*  
*Somatochlora hineana*



*Somatochlora  
albicincta*




*Somatochlora  
cingulata*







*Somatochlora  
whitehousei*

**CORDULIIDAE 5    *Somatochlora albicincta, cingulata, whitehousei***



## CORDULIIDAE 6







*Somatochlora brevicincta, sahlbergi, septentrionalis,  
hudsonica, semicircularis, williamsoni*




## CORDULIIDAE 7

*Somatochlora kennedyi, calverti, margarita, forcipata, franklinii*









Source: Donnelly, T.W. 2008. A distributional synthesis of dragonflies (Odonata) from the Americas. Part 7. Subfamily Corduliidae. *Entomological News* 120(1): 1–100.



**CORDULIIDAE 8**    *Somatochlora elongata, ensigera, filosa, linearis,  
georgiana, ozarkensis*











**CORDULIIDAE 9** *Somatochlora provocans, tenebrosa, minor, walshii*











## CORDULIIDAE 10; LIBELLULIDAE 1

*Somatochlora incurvata*  
*Brachymesia furcata, herbida*  
*Brechmorhogax mendax, pertinax*  
*Macrothemis inacuta*  
*Cannaphila insularis*




## LIBELLULIDAE 2

*Brachymesia gravida*  
*Macrothemis imitans, inequiunguis*  
*Macrodiplax balteata*  
*Dythemis fugax, maya, nigrescens, velox*












### LIBELLULIDAE 3

*Idiataphe cubensis*  
*Paltothemis lineatipes*  
*Micrathyria aequalis, didyma, hagenii*  
*Erythrodiplax funerea, minuscula*  
*Tholymis citrina*




#### LIBELLULIDAE 4

*Erythemis simplicicollis, attala, peruviana*  
*Erythrodiplax basifusca, fusca*  
*Tauriphila australis, azteca*




**LIBELLULIDAE 5**


*Erythrodiplax umbrata*  
*Erythemis vesiculosa, plebeja*  
*Tramea insularis, binotata*  
*Pseudoleon superbus*  
*Miathyria marcella*  
*Celithemis amanda, bertha*




*Tramea  
calverti*




*Ladona  
deplanata*




*Tramea  
carolina*




*Ladona  
exusta*



*Tramea  
abdominalis*




*Celithemis  
verna*




*Celithemis  
martha*


**LIBELLULIDAE 6**    *Ladona deplanata, exusta*  
                         *Tramea calverti, carolina,*  
                         *abdominalis*  
                         *Celithemis verna, martha*




*Celithemis  
elisa*




*Celithemis  
eponina*




*Celithemis  
ornata*



*Celithemis  
fasciata*



*Ladona  
julia*








*Leucorrhinia  
frigida*

## LIBELLULIDAE 7

*Celithemis elisa, ornata, eponina, fasciata*


*Celithemis ornata*

*Leucorrhinia frigida*









## LIBELLULIDAE 8

*Leucorrhinia proxima, glacialis, hudsonica*  
*Libellula forensis*  
*Sympetrum illotum*




## LIBELLULIDAE 9


*Leucorrhinia patricia, intacta, borealis*  
*Libellula nodisticta, flava, Jesseana*




**LIBELLULIDAE 10    *Libellula auripennis, axilena, comanche, croceipennis, cyanea, needhami***




*Libellula  
incesta*




*Libellula  
vibrans*




*Libellula  
composita*



*Libellula  
saturata*




*Libellula  
semifasciata*



*Nannothemis  
bella*


## LIBELLULIDAE 11

*Libellula incesta, vibrans, composita,  
saturata, semifasciata  
Nannothemis bella*



**LIBELLULIDAE 12**

*Libellula pulchella, luctuosa, quadrimaculata*




### LIBELLULIDAE 13


*Orthemis ferruginea, discolor, antillean species*

*Perithemis domitia, intensa, tenera*




*Sympetrum signiferum*



**LIBELLULIDAE 14** *Pachydiplax longipennis; Pantala flavescens, hymenaea*




## LIBELLULIDAE 15




**LIBELLULIDAE 16**

*Sympetrum semicinctum, danae, costiferum*



**LIBELLULIDAE 17** *Sympetrum corruptum, internum, obtrusum (incl. hybrid)*



**LIBELLULIDAE 18** *Tramea lacerata, onusta*  
*Sympetrum vicinum*